- Tell your doctor if you ever had a severe reaction after a dose of influenza vaccine.
- Tell your doctor if you ever had Guillain-Barré Syndrome (a severe paralytic illness, also called GBS). Your doctor will help you decide whether the vaccine is recommended for you.
- People who are moderately or severely ill should usually wait until they recover before getting flu vaccine. If you are ill, talk to your doctor about whether to reschedule the vaccination. People with a mild illness can usually get the vaccine.

5 What are the risks from inactivated influenza vaccine?

A vaccine, like any medicine, could possibly cause serious problems, such as severe allergic reactions. The risk of a vaccine causing serious harm, or death, is extremely small.

Serious problems from inactivated influenza vaccine are very rare. The viruses in inactivated influenza vaccine have been killed, so you cannot get influenza from the vaccine.

Mild problems:
- soreness, redness, or swelling where the shot was given
- hoarseness; sore, red or itchy eyes; cough
- fever • aches • headache • itching • fatigue

If these problems occur, they usually begin soon after the shot and last 1-2 days.

Moderate problems:
Young children who get inactivated flu vaccine and pneumococcal vaccine (PCV13) at the same time appear to be at increased risk for seizures caused by fever. Ask your doctor for more information.

Tell your doctor if a child who is getting flu vaccine has ever had a seizure.

Severe problems:
- Life-threatening allergic reactions from vaccines are very rare. If they do occur, it is usually within a few minutes to a few hours after the shot.
- In 1976, a type of inactivated influenza (swine flu) vaccine was associated with Guillain-Barré Syndrome (GBS). Since then, flu vaccines have not been clearly linked to GBS. However, if there is a risk of GBS from current flu vaccines, it would be no more than 1 or 2 cases per million people vaccinated. This is much lower than the risk of severe influenza, which can be prevented by vaccination.

The safety of vaccines is always being monitored. For more information, visit:
www.cdc.gov/vaccinesafety/Vaccine_Monitoring/Index.html and **www.cdc.gov/vaccinesafety/Activities/Activities_Index.html**

One brand of inactivated flu vaccine, called Afluria, **should not be given** to children 8 years of age or younger, except in special circumstances. A related vaccine was associated with fevers and fever-related seizures in young children in Australia. Your doctor can give you more information.

6 What if there is a severe reaction?

What should I look for?
- Any unusual condition, such as a high fever or unusual behavior. Signs of a serious allergic reaction can include difficulty breathing, hoarseness or wheezing, hives, paleness, weakness, a fast heart beat or dizziness.

What should I do?
- **Call** a doctor, or get the person to a doctor right away.
- **Tell** your doctor what happened, the date and time it happened, and when the vaccination was given.
- **Ask** your doctor, nurse, or health department to report the reaction by filing a Vaccine Adverse Event Reporting System (VAERS) form. Or you can file this report through the VAERS web site at **www.vaers.hhs.gov**, or by calling **1-800-822-7967**.

VAERS does not provide medical advice.

7 The National Vaccine Injury Compensation Program

The National Vaccine Injury Compensation Program (VICP) was created in 1986.

People who believe they may have been injured by a vaccine can learn about the program and about filing a claim by calling **1-800-338-2382** or visiting the VICP website at **www.hrsa.gov/vaccinecompensation**.

8 How can I learn more?

- Ask your doctor. They can give you the vaccine package insert or suggest other sources of information.
- Call your local or state health department.
- Contact the Centers for Disease Control and Prevention (CDC):
  - Call **1-800-232-4636 (1-800-CDC-INFO)** or
  - Visit CDC's website at **www.cdc.gov/flu**

Office Use Only